arXiv:1606.00723v2 [math.DS] 23 Jun 2016

On deformation spaces of quadratic rational functions

Tanya Firsova[1], *Jeremy Kahn* [2], *Nikita Selinger*

Abstract

We study the group of self-equivalences of a partially postcritically finite branched cover and answer a question of Adam Epstein about contractibility and connectedness of certain deformation spaces of rational maps.

1. Introduction

A celebrated theorem of W. Thurston provides a topological criterion for when a postcritically finite branched covering is equivalent to a rational map. The cornerstone of Thurston's classification is the pull-back map on the Teichmüller space induced by the branched cover.

Let P_f be the postcritical set of a branched cover f; consider the Teichmüller space $\mathrm{Teich}(\mathbb{S}^2, P_f)$ of the topological 2-sphere $\mathbb{S}^2$ with the set P_f marked. Then by taking the composition of the pull-back $\tilde{f^*} : \mathrm{Teich}(\mathbb{S}^2, P_f) \to \mathrm{Teich}(\mathbb{S}^2, f^{-1}(P_f))$ with the forgetful map $\tilde{i} : \mathrm{Teich}(\mathbb{S}^2, f^{-1}(P_f)) \to \mathrm{Teich}(\mathbb{S}^2, P_f)$, one obtains the Thurston pull-back map

$$f^* : \mathrm{Teich}(\mathbb{S}^2, P_f) \to \mathrm{Teich}(\mathbb{S}^2, P_f).$$

The map f^* has a fixed point in Teichmüller space if and only if f equivalent to a rational map.

Now assume $A \subset B \subset \mathbb{S}^2$ are finite sets, $f : \mathbb{S}^2 \to \mathbb{S}^2$ is a branched cover, $V(f)$ is the set of critical values of f. Moreover, assume that

$$f(A) \cup V(f) \subset B.$$

Then there is a well-defined map $f^* : \mathrm{Teich}(\mathbb{S}^2, B) \to \mathrm{Teich}(\mathbb{S}^2, A)$, defined by pulling back by f and then erasing $f^{-1}(B) \backslash A$.

Following Adam Epstein, we define the deformation space $\mathrm{Def}(f, A, B)$:

$$\mathrm{Def}(f, A, B) := \{\tau \in \mathrm{Teich}(\mathbb{S}^2, B) : \quad f^*\tau = i_*\tau\},$$

where $f^* : \mathrm{Teich}(\mathbb{S}^2, B) \to \mathrm{Teich}(\mathbb{S}^2, A)$ is the pull-back map defined above, $i_* : \mathrm{Teich}(\mathbb{S}^2, B) \to \mathrm{Teich}(\mathbb{S}^2, A)$ is the forgetful map. The deformation space is non-empty if and only if f is equivalent to a rational map, that is, there exist a rational map $g \colon \hat{\mathbb{C}} \to \hat{\mathbb{C}}$ and $\phi, \psi \colon \mathbb{S}^2 \to \hat{\mathbb{C}}$ such that $\phi \sim \psi$ rel A and the following diagram commutes:

$$\begin{array}{ccc} (\mathbb{S}^2, A) & \xrightarrow{\psi} & (\hat{\mathbb{C}}, \psi(A)) \\ \big\downarrow f & & \big\downarrow g \\ (\mathbb{S}^2, B) & \xrightarrow{\phi} & (\hat{\mathbb{C}}, \phi(B)). \end{array}$$

[1]Research of the first author was supported in part by NSF.

[2]Research of the second author was supported in part by NSF.

In fact, each structure $\tau \in \mathrm{Def}(f, A, B)$ defines such a rational map up to conjugacy by Möbius transformations.

Adam Epstein proved [2] that in non-exceptional cases the dimension of $\mathrm{Def}(f, A, B)$ is equal to $\#(B \setminus A)$ and asked a question whether deformation spaces are contractible. Below we give a simple condition that guarantees that the deformation space is not contractible. We prove that deformation spaces corresponding to maps in Per_n, $n \geq 4$ satisfy this condition. Together with results of Mary Rees (see Theorem 3.4 below) this implies that each of these spaces consists of countably many connected components. The last statement has been shown for maps in Per_4 independently by Eriko Hironaka and Sarah Koch [3].

1.1. **Acknowledgements.** We would like to thank Jan Kiwi for his valuable suggestions.

2. Background. Notations.

Let $P \subset \mathbb{S}^2$ be finite. We will use the following notations:

- $\mathrm{Teich}(P)$ is the Teichmüller space of the sphere with the marked set P,
- $\mathrm{Mod}(P)$ is the moduli space of the sphere with the marked set P,
- $\mathrm{MCG}(P)$ is the (pure) mapping class group of $\mathrm{Teich}(P)$,
- $\pi : \mathrm{Teich}(P) \to \mathrm{Mod}(P)$ is the natural projection from the Teichmüller space $\mathrm{Teich}(P)$ to the moduli space $\mathrm{Mod}(P)$.

In our setting, the Teichmüller space is the universal cover of the moduli space. The mapping class group $\mathrm{MCG}(P)$ is naturally identified with the group of deck transformations of the cover π.

Recall that an element of $\mathrm{Teich}(B)$ is an equivalence class of homeomorphisms $\phi : (\mathbb{S}^2, B) \to (\hat{\mathbb{C}}, \phi(B))$. Two such homeomorphisms ϕ_1 and ϕ_2 are equivalent if there exist a homeomorphism $\beta : (\mathbb{S}^2, P) \to (\mathbb{S}^2, P)$, homotopic to the identity, and a Möbius transformation α, such that the following diagram commutes.

$$\begin{array}{ccc} (\mathbb{S}^2, P) & \xrightarrow{\phi_1} & (\hat{\mathbb{C}}, \phi_1(P)) \\ \downarrow \beta & & \downarrow \alpha \\ (\mathbb{S}^2, P) & \xrightarrow{\phi_2} & (\hat{\mathbb{C}}, \phi_2(P)) \end{array}$$

Consider non-trivial, non-peripheral, simple closed curves on $(\mathbb{S}^2 \setminus P)$. Let $\mathcal{C}(P)$ be the set of equivalence classes of such curves up to free homotopy relative to P. We say that two distinct equivalence classes do not intersect if they have disjoint representatives. We denote the class of $\gamma \in (\mathbb{S}^2, P)$ by $[\gamma]_P$. We say that $\Gamma = \{\gamma_1, \ldots, \gamma_n\}$ is *a multicurve* on $(\mathbb{S}^2, P)$ if the curves γ_i represent mutually disjoint equivalence classes. Two multicurves Γ, $\tilde{\Gamma}$ are equivalent if for each $\gamma \in \Gamma$ there exists $\tilde{\gamma} \in \tilde{\Gamma}$ such that $[\gamma]_P = [\tilde{\gamma}]_P$ and vice versa. We denote by $\mathcal{M}(P)$ the set of equivalence classes of multicurves.

Let Γ be a multicurve on $(\mathbb{S}^2, P)$; we denote by $(\mathbb{S}^2, P)/\Gamma$ the topological space with marked points obtained by collapsing each curve in Γ to a point. This space, a nodal sphere, is a collection of spheres with marked points attached to each other at special points that we will call *nodes*. Note that since every curve in Γ is non-peripheral and no two curves in Γ are homotopic to each other, there are at least three special points (either marked points, or nodes) on every sphere in the collection.

We will call the Teichmüller space of the nodal sphere $(\mathbb{S}^2, P)/\Gamma$, defined as the product of Teichmüller spaces of components of that sphere, *the stratum* $\mathcal{S}_\Gamma$ corresponding to the (equivalence class of) multicurve Γ. Taking the union of all strata we obtain the augmented Teichmüller space:

$$\overline{\mathrm{Teich}}(P) := \bigcup_{\Gamma \in \mathcal{M}(P)} \mathcal{S}_\Gamma.$$

Note that, by definition, $\mathrm{Teich}(P) = \mathcal{S}_\emptyset \subset \overline{\mathrm{Teich}}(P)$.

A point on a stratum $\mathcal{S}_\Gamma$ can be represented by a continuous map ϕ from $(\mathbb{S}^2, P)$ onto a nodal Riemann surface with marked points which is surjective and injective except that each curve in Γ is collapsed to a node. The equivalence relation on such maps is a straightforward generalization of the equivalence relation on homeomorphisms defined above.

The geodesic length functions l_γ for $\gamma \in \mathcal{C}(P)$ extend to $\overline{\mathrm{Teich}}(P)$ by setting $l_\gamma = 0$ if γ is homotopic to a node, $l_\gamma = \infty$ if γ has non-trivial intersection with at least one of the nodes. In the remaning case, γ is a curve on one of the components of the nodal surface and we measure l_γ with respect to the hyperbolic metric of that component. The topology of the augmented Teichmüller space can be defined by requiring that $\tau_k \to \tau$ if and only if $l_\gamma(\tau_k) \to l_\gamma(\tau)$ for all $\gamma \in \mathcal{C}(P)$.

Theorem 2.1. *The map f^* extends continuously to the augmented Teichmüller space.*

Proof. The proof of Theorem 6.4 in [7] applies mutatis mutandis to our setting. □

We say that a triple (f, A, B) and, correspondingly, the deformation space $\mathrm{Def}(f, A, B)$ are *reduced* if

$$B = A \cup f(A) \cup V(f).$$

Below we assume that deformation spaces $\mathrm{Def}(f, A, B)$ we consider are reduced.

Theorem 2.2. [2] *If f is not exceptional, i.e. f is not a flexible Lattes example, then* $\mathrm{Def}(f, A, B)$ *is either empty or a smooth submanifold of* $\mathrm{Teich}(B)$ *of dimension $\#B - \#A$.*

Throughout the article, we consider only non-exceptional branched covers.

3. Generalities about Deformation spaces

3.1. **Equivalences, geometric equivalences, and self-equivalences.** We are going to use $\mathbb{S}^2$ and $\hat{\mathbb{C}}$ in a way where they are more or less equivalent, and yet distinct. We could think of $\hat{\mathbb{C}}$ as $\mathbb{S}^2$ with the standard complex structure. So from this point of view $\mathbb{S}^2$ is the image of $\hat{\mathbb{C}}$ by the forgetful functor from Riemann surfaces to smooth (or topological) surfaces.

Now, suppose, for $i = 0, 1$, that $f_i \colon (\mathbb{S}^2, A_i) \to (\mathbb{S}^2, B_i)$ are branched covers, with $A_i \cup V(f_i) \subset B_i$. We say that a homeomorphism $h \colon (\mathbb{S}^2, A_0, B_0) \to (\mathbb{S}^2, A_1, B_1)$ is an *equivalence* (or *topological equivalence*) if there is a $h' \colon (\mathbb{S}^2, A_0) \to (\mathbb{S}^2, A_1)$ such that $h \circ f_0 = f_1 \circ h'$, and $h' \sim h$ rel A_0. We observe that this is Thurston's notion of equivalence in the case where $A_i = B_i$.

Now replace $\mathbb{S}^2$ with $\hat{\mathbb{C}}$ in the definition in the previous paragraph. We do not suppose that the f_i are holomorphic. We say that an equivalence $h \colon (\hat{\mathbb{C}}, A_0, B_0) \to (\hat{\mathbb{C}}, A_1, B_1)$ is *geometric* if it is conformal (we do not require h' to be conformal).

We define the *Rees component* $\mathcal{R}(f)$ as the space of branched covers $g \colon (\hat{\mathbb{C}}, A') \to (\hat{\mathbb{C}}, B')$ that are topologically equivalent to f, modulo geometric equivalence. The space $\mathcal{R}(f)$ is a connected component of the space of all brached covers with the same critical portrait as f, modulo the same equivalence relation. We let $\mathcal{R}_{\mathrm{Rat}}(f, A, B) \subset \mathcal{R}(f)$ be the equivalence classes with holomorphic representatives. We have the following natural map

$$\pi_{\mathcal{TR}} \colon \mathrm{Teich}(B) \to \mathcal{R}(f).$$

For $\tau \in \mathrm{Teich}(B)$, let $\phi \colon (S^2, B) \to (\hat{\mathbb{C}}, \phi(B))$ be a representative, then we define $\pi_{\mathcal{TR}}(\tau) = \phi \circ f \circ \phi^{-1}$. Choosing a different representative ϕ' of the structure τ, we obtain the map $\phi' \circ f \circ (\phi')^{-1}$ which is geometrically equivalent to $\phi \circ f \circ \phi^{-1}$; we see that the map $\pi_{\mathcal{TR}}$ is well-defined.

The map $\pi_{\mathcal{TR}}$ is surjective. Let $g \colon (\hat{\mathbb{C}}, A') \to (\hat{\mathbb{C}}, B')$ be a branched cover topologically equivalent to f, then there exists $\phi \colon (\mathbb{S}^2, B) \to (\hat{\mathbb{C}}, B')$, $\phi' \colon (\mathbb{S}^2, A) \to (\hat{\mathbb{C}}, A')$ such that $g \circ \phi' = \phi \circ f$, $\phi \sim \phi'$ rel A. The branched cover g is thus geometrically equivalent to $\phi \circ f \circ \phi^{-1}$.

Moreover, if $h \in \mathrm{MCG}(S^2, B)$ is a self-equivalence of f, then $\pi_{\mathcal{TR}}(h\tau) = \pi_{\mathcal{TR}}(\tau)$. The map $\pi_{\mathcal{TR}} \colon \mathrm{Teich}(B) \to \mathcal{R}(f)$ is a covering map. It is a Galois cover, since $\mathrm{Teich}(B)$ is simply connected. We can identify $G_f := \mathrm{Deck}(\mathrm{Teich}(B)/\mathcal{R}(f))$ with the group of homotopy classes of self-equivalences of (f, A, B).

We also have the following map:

$$\pi_{\mathcal{RM}} \colon \mathcal{R}(f) \to \mathrm{Mod}(B)$$

where $\pi_{\mathcal{RM}}(f', A', B') := B'$. The map $\pi_{\mathcal{RM}}$ is a covering and

$$\pi = \pi_{\mathcal{RM}} \circ \pi_{\mathcal{TR}}.$$

We let $\mathcal{R}_{\mathrm{Rat}}(f,A,B) \subset \mathcal{R}(f)$ be the geometric equivalence classes that are represented by rational maps (holomorphic branched covers).

Note that the group of self-equivalences G_f and the Rees component $\mathcal{R}(f)$ are well-defined even if the branched cover f is not realized by a rational function. Now we assume that f can be realized by a rational function, i.e. $\mathcal{R}_{\mathrm{Rat}}(f,A,B)$ and $\mathrm{Def}(f,A,B)$ are non-empty.

Lemma 3.1. *The map $\pi_{\mathcal{TR}}\colon \mathrm{Def}(f,A,B) \to \mathcal{R}_{\mathrm{Rat}}(f,A,B)$ is a Galois covering map.*

Proof. Let $\tau \in \mathrm{Def}(f,A,B)$; let $\phi : (\mathbb{S}^2, B) \to (\hat{\mathbb{C}}, \phi(B))$ be a representative of τ. Then by definition of the deformation space, there exists $\phi' : (\mathbb{S}^2, A) \to (\hat{\mathbb{C}}, \phi(A))$, $\phi' \sim \phi$ rel A, and a rational map $g\colon (\hat{\mathbb{C}}, \phi(A)) \to (\hat{\mathbb{C}}, \phi(B))$ such that $g \circ \phi' = \phi \circ f$. Let h be a self-equivalence and $\tau' \in \mathrm{Teich}(B)$ a structure defined by $\phi \circ h$. We have the following diagram:

$$\begin{array}{ccccc}
(\mathbb{S}^2, A) & \xrightarrow{h'} & (S^2, A) & \xrightarrow{\phi'} & (\hat{\mathbb{C}}, \phi(A)) \\
\big\downarrow f & & \big\downarrow f & & \big\downarrow g \\
(\mathbb{S}^2, B) & \xrightarrow{h} & (S^2, B) & \xrightarrow{\phi} & (\hat{\mathbb{C}}, \phi(B))
\end{array}$$

Hence, $\tau' \in \mathrm{Def}(f,A,B)$. So, the group G_f acts on $\mathrm{Def}(f,A,B)$, hence $\mathcal{R}_{\mathrm{Rat}}(f,A,B)$ is an image of $\mathrm{Def}(f,A,B)$ under $\pi_{\mathcal{TR}}$. □

We say that an element $h \in \mathrm{MCG}(B)$ is *liftable* if there exists $h' \in \mathrm{MCG}(A)$ and their representatives $\tilde{h}$ and $\tilde{h}'$ such that $f \circ \tilde{h} = \tilde{h}' \circ f$. If such an element $h' \in \mathrm{MCG}(A)$ exists, then it is unique. Hence, f acts on liftable elements, we will use the notation $f^*\colon H_f \to \mathrm{MCG}(A)$ for the action $h \mapsto h'$. The group of liftable elements $H_f < \mathrm{MCG}(B)$ is a finite index subgroup [6]. The group G_f is a subgroup of H_f.

(1) $\mathcal{R}_{\mathrm{Rat}}(f,A,B) = \pi_{\mathcal{TR}}(\mathrm{Def}(f,A,B))$.
Since restriction of $\pi_{\mathcal{TR}}$ to $\mathrm{Def}(f,A,B)$ is a Galois covering, $\mathcal{R}_{\mathrm{Rat}}(f,A,B)$ is a smooth manifold. A priori, it can consist of several components.

(2) Let $M(f,A,B) = \pi_{RM}(\mathcal{R}_{\mathrm{Rat}}(f,A,B))$. We can also consider the space $\mathrm{Rat}(f,A,B)$ of all maps with the same critical portrait as f. We define $\mathrm{Rat}(f,A,B)$ as the space of pairs $g\colon \hat{\mathbb{C}} \to \hat{\mathbb{C}}, h\colon B \to \hat{\mathbb{C}}$ such that g is holomorphic, $g \circ h = h \circ f$ on A, and the local degree of f at x equals to the local degree of g at $h(x)$ for all $x \in A$. We have a natural embedding of $\mathcal{R}_{\mathrm{Rat}}(f,A,B)$ into $\mathrm{Rat}(f,A,B)$. Hence, the restriction of $\pi_{\mathcal{RM}}$ to $\mathcal{R}_{\mathrm{Rat}}(f,A,B)$ is necessarily a finite degree covering map. If it is one-to-one, we say that the deformation space is *projectable*.

We summarize the notations in the following diagram:



We say that a topological space embeds into another topological space *injectively on* π_1 if the induced homeomorphism on fundamental groups is injective.

Rees in [5] considered the space $\mathcal{B}(f, A, B)$ – the connected component containing f of the space of all branched covers topologically equivalent to f, modulo conjugation by Möbius transformations. She studied quadratic rational maps with a critical cycle and showed [5, Section 5.10] the following (see Section 3.3 for the definition of Per'_n).

Theorem 3.2. *For $f \in \mathrm{Per}'_n$, the space $\mathrm{Rat}(f, A, B)$ embeds injectively on π_1 into $\mathcal{B}(f, A, B)$.*

The space $\mathcal{R}(f)$ is the quotient of $\mathcal{B}(f, A, B)$ by the following equivalence relation: $(f, A, B) \equiv (g, A, B)$ if and only if f is homotopic to g relative to A.

Corollary 3.3. *For $f \in \mathrm{Per}'_n$, the space $\mathrm{Rat}(f, A, B)$ embeds injectively on π_1 into $\mathcal{R}(f)$.*

Proof. We observe that $\mathcal{B}(f, A, B)$ is a fibration over $\mathcal{R}(f)$ where a fiber over (f', A', B') is the set of all homeomorphisms that are homotopic to the identity relative to A'. Since every fiber is contractible, $\mathcal{R}(f)$ is a deformation retract of $\mathcal{B}(f, A, B)$. □

The previous two statements immediately imply the following.

Theorem 3.4. *For $f \in \mathrm{Per}'_n$ each component of $\mathrm{Def}(f, A, B)$ is contractible.*

Proof. The embedding of $\mathcal{R}_{\mathrm{Rat}}(f, A, B)$ to $\mathcal{R}(f)$ is injective on π_1 if and only if each component of $\mathrm{Def}(f, A, B)$ is contractible. Combining Corollary 3.3 and Theorem 3.2 we get the desired result. □

3.2. **Equalizing curves.** The forgetful map induces the map $i_* : \mathcal{C}(B) \to \mathcal{C}(A)$ on classes of simple closed curves:

$$i_* : [\gamma]_B \mapsto [\gamma]_A.$$

Similarly, we define the action of $i_* : \mathcal{M}(B) \to \mathcal{M}(A)$.

A formal linear sum

$$m_1\gamma_1 + \cdots + m_n\gamma_n,$$

where $m_i \in \mathbb{Q}$ is called a weighted multicurve. Let $\mathcal{W}(P)$ be a space of weighted multicurves on $(\mathbb{S}^2, P)$. The forgetful map extends linearly to $i_*\colon \mathcal{W}(B) \to \mathcal{W}(A)$.

Let $\beta_1, \dots, \beta_k \subset (\mathbb{S}^2, f^{-1}(B))$ be preimages of a simple closed curve γ under the map f; denote by d_i the degree of f restricted to β_i. Define

$$\tilde{f}^*(\gamma) = \sum_{i=1}^{k} \frac{1}{d_i}\beta_i.$$

We extend this map linearly to $\tilde{f}^* : \mathcal{W}(B) \to \mathcal{W}(f^{-1}(B))$. Let

$$f^* := \tilde{i}_* \circ \tilde{f}^*,$$

where $\tilde{i}_*$ is the forgetful map $\tilde{i}_* : \mathcal{W}(f^{-1}(B)) \to \mathcal{W}(A)$.

We also get a well-defined map

$$f^* : \mathcal{M}(P) \to \mathcal{M}(P),$$

if we ignore the weights in the definition of f^*.

To a pair of multicurves $\Gamma = \{\gamma_i\}$ on $(\mathbb{S}^2, B)$ and $\Delta = \{\delta_j\}$ on $(\mathbb{S}^2, A)$, we associate the Thurston matrix T_Γ^Δ which is obtained from the map $f^* : \mathcal{W}(B) \to \mathcal{W}(A)$ by restricting the domain to $\mathbb{Q}^\Gamma$ and projecting the result to $\mathbb{Q}^\Delta$. The (i,j)-th entry of T_Γ^Δ equals to $\sum \frac{1}{d_{i,j,\alpha}}$, where $d_{i,j,\alpha}$ is the degree of f restricted to $\delta_{j,\alpha}$ and $\{\delta_{j,\alpha}\}$ is the set of connected components of $f^{-1}(\gamma_i)$ that are homotopic to δ_j. In a similar fashion we define the restriction of i_*, the matrix I_Γ^Δ, where (i,j)-th entry is 1 if $[\gamma_i]_A = [\delta_j]_A$ and 0 otherwise.

We say that a multicurve Γ is *equalizing* if $f^*\Gamma = i_*\Gamma$ as multicurves. A multicurve $\Gamma = \{\gamma_1, \dots, \gamma_n\}$ is *arithmetically equalizing* if there exists a set of positive rational weights $\{m_1, \dots, m_n\}$ such that $i_*[m_1\gamma_1 \cdots + m_n\gamma_n]_B = f^*[m_1\gamma_1 + \cdots + m_n\gamma_n]_B$.

Theorem 3.5. *$f^*\mathcal{S}_\Gamma = i_*\mathcal{S}_\Gamma$ if and only if Γ is an equalizing multicurve. If the deformation space accumulates on $\mathcal{S}_\Gamma$, then there exists $\tilde{\Gamma} \subset \Gamma$ such that $\tilde{\Gamma}$ is an arithmetically equalizing multicurve.*

Proof. By Theorem 2.1, the pull-back map f^* extends continuously to the augmented Teichmüller space and $f^*(S_\Gamma) = S_{f^*(\Gamma)}$. The forgetful map extends continuously to the augmented Teichmüller space as well. Hence, $i_*(S_\Gamma) = S_{i_*(\Gamma)}$ and the first statement follows.

Let $\tau^k \in \mathrm{Def}(f, A, B)$ be a sequence of conformal structures such that $\tau^k \to \tau^*$, $\tau^* \in \mathcal{S}_\Gamma$. Let m_i^k be the modulus of the maximal embedded annulus homotopic to γ_i in the Riemann sphere corresponding to τ^k. Set $\Delta = f^{-1}(\Gamma)$; Let n_j^k be the modulus of the maximal embedded annulus homotopic to δ_j in the Riemann sphere corresponding to $f^*(\tau^k)$. Applying Theorem 7.1 from [1], we see that $n^k = T_\Gamma^\Delta m^k + O(1)$ where the additive constant depends only on f and the supremum of conformal moduli of curves not in Γ. Similarly, we get $n^k = I_\Gamma^\Delta m^k + O(1)$ because $f^*(\tau) = i_*(\tau)$ by definition of $\mathrm{Def}(f, A, B)$. Hence, $(T_\Gamma^\Delta - I_\Gamma^\Delta)m^k = O(1)$.

Since all entries of m^k tend to infinity, we see that there is a non-negative rational solution to $(T_\Gamma^\Delta - I_\Gamma^\Delta)m = 0$. □

In Lemma 5.3 we give an example of a curve that is arithmetically equalizing, but the deformation space does not accumulate to the corresponding stratum.

For a simple curve γ, we denote by T_γ a Dehn twist along γ. Suppose that m is such that, for every curve β that maps to γ by f, the degree of f on β divides m. Then $f^*(m\gamma) = \sum n_i\beta_i$, where the n_i are integers, and $f^*(T_\gamma^m) = \prod T_{\beta_i}^{n_i}$. In particular, $T_\gamma^{d!}$ is always liftable, for every γ, where d is the degree of f.

Lemma 3.6. *Let $\Gamma = \{\gamma_i\}$ be an arithmetically equalizing multicurve. Assume $f^*(m_1\gamma_1 + \cdots + m_n\gamma_n) = i_*(m_1\gamma_1 + \dots m_n\gamma_n)$, where $m_i \in \mathbb{Z}$. Moreover, assume that m_i are such that, $T_{\gamma_i}^{m_i}$ are liftable. Then $T_{\gamma_1}^{m_1} \circ \cdots \circ T_{\gamma_n}^{m_n} \in G_f$.*

Proof. Let $i_*(m_1\gamma_1 + \dots m_n\gamma_n) = l_1\delta_1 + \cdots + l_k\delta_k$, where δ_i are different simple non-peripheral curves in $(\mathbb{S}^2, A)$. Then $i_*\left(T_{\gamma_1}^{m_1} \circ \cdots \circ T_{\gamma_n}^{m_n}\right) = f^*\left(T_{\gamma_1}^{m_1} \circ \cdots \circ T_{\gamma_n}^{m_n}\right) = T_{\delta_1}^{l_1} \circ \cdots \circ T_{\delta_k}^{l_k}$. □

Corollary 3.7. *If $\mathrm{Def}(f, A, B)$ accumulates to a point $p \in \mathcal{S}_\Gamma$, $\Gamma = \{\gamma_1, \dots, \gamma_n\}$, then there exist $m_1, \dots, m_n \in \mathbb{Z}$ such that $T_{\gamma_n}^{m_1} \circ \cdots \circ T_{\gamma_n}^{m_n} \in G_f$.*

Proof. The corollary is a direct consequence of Theorem 3.5 and Lemma 3.6 □

3.3. **Quadratic rational map.** Let $P(z)$, $Q(z)$ be polynomials with no common roots. The map $f(z) = \frac{P(z)}{Q(z)}$ is called a degree $d := \max\{\deg P, \deg Q\}$ rational map. The space of rational maps of degree d is denoted by Rat_d. Specifying dynamical relations for orbits of critical points, one obtains subvarieties of Rat_d with prescribed dynamical properties. Möbius transformations act on the space of rational maps by conjugating. The space of conjugacy classes of rational maps of degree 2 with marked critical points is denoted by $\mathcal{M}_2^{cm}$. Milnor [4] showed that $\mathcal{M}_2^{cm}$ is biholomorphic to $\mathbb{C}^2$.

Quadratic maps have two critical points. We denote them by c_1 and c_2. We will use the notation $v = f(c_2)$. Consider the variety $\mathrm{Per}_n \subset \mathcal{M}_2^{cm}$:

$$\mathrm{Per}_n := \{f \in \mathcal{M}_2^{cm} : \ f^n(c_1) = c_1 \text{ and } f^k(c_1) \neq c_1 \text{ when } 0 < k < n\}.$$

Per_n is an algebraic subvariety of $\mathcal{M}_2^{cm}$. It is not known whether Per_n is irreducible for all n.

Suppose that $c_1, f(c_1)$, and $f^2(c_1)$ are mutually distinct for a map $f \in \mathcal{M}_2^{cm}$. Conjugating by a Möbius transformations, we may assume that $c_1 = 0$ and $0 \mapsto \infty \mapsto 1$. With this parametrization f takes the form

$$z \mapsto 1 + \frac{b}{z} + \frac{c}{z^2} =: f_{b,c}(z).$$

On the other hand, any two values of $b \in \mathbb{C}$, $c \in \mathbb{C}^*$ define a rational map in $\mathcal{M}_2^{cm}$.

Let

$$\mathrm{Per}_n' = \{f \in \mathrm{Per}_n : \ v \neq f^k(c_1), k = 0, 1, \dots\}$$

Let $f \in \mathrm{Per}_n'$. Take $A = \{c_1, f(c_1) \dots, f^{(n-1)}(c_1)\}$, $B = A \cup \{v\}$. Then $\mathrm{Def}(f, A, B)$ is 1-dimensional.

Lemma 3.8. *Let $f \in \mathrm{Per}_3$, $A = \{0, 1, \infty\}$, $B = A \cup \{v\}$. The deformation space $\mathrm{Def}(f, A, B)$ is not projectable. The map $\pi_{\mathcal{RM}}|_{\mathcal{R}_{\mathrm{Rat}(f)}}$ is a two-to-one covering.*

Proof. Recall that with our standard parametrization, a map $f \in \mathcal{M}_2^{cm}$ has the form

$$f_{b,c}(z) = 1 + \frac{b}{z} + \frac{c}{z^2}.$$

If $f_{b,c} \in \mathrm{Per}_3$, $f(1) = 1 + b + c = 0$, so $c = 1 - b$. We compute the second critical value $v = 1 - \frac{b^2}{4(1-b)}$. The last equation has two solutions b_1, b_2 for a fixed v, so the maps $f_{b_1,1-b_1}$ and $f_{b_2,1-b_2}$ project to the same point in the moduli space. □

Lemma 3.9. *Let $f \in \mathrm{Per}_n$ be such that $v \neq f^k(c_1)$ for any $k \in \mathbb{N}$. If $n \geq 4$, the deformation space $\mathrm{Def}(f, A, B)$ is projectable.*

Proof. Let $\rho = f(1)$, $s = f(\rho)$. Then we have the following system of linear equations:

$$1 + b + c = \rho$$
$$\rho^2 + b\rho + c = \rho^2 s$$

If $n \geq 4$, $s \neq \infty$, so the system of linear equations is well-defined. Its determinant is equal to $(1 - \rho) \neq 0$. Therefore, the values of ρ and s uniquely determine the map. □

4. Topology of the Deformation space

In this section we assume that $\mathrm{Def}(f, A, B)$ is projectable. The space $M(f, A, B)$ can a prioi contain several components. Let M_0 be one of the components. Denote by $I \colon M_0 \to \mathrm{Mod}(B)$ the inclusion map. Fix a point $m_0 \in M_0$; the inclusion map induces a homomorphism $I_* \colon \pi_1(M_0, m_0) \to \pi_1(\mathrm{Mod}(B), m_0) = \mathrm{MCG}(B)$.

$G_0 := I_*(\pi_1(M_0, m_0)) \cong \pi_1(M_0, m_0)/\ker I_*$ is a subgroup of G_f; it is the group of self-equivalences that can be realized by paths on M_0.

Let Def_0 be a component of the deformation space $\mathrm{Def}(f, A, B)$ such that $\pi(\mathrm{Def}(f, A, B)) = M_0$. It follows that $\pi_1(\mathrm{Def}_0) = \ker I_*$; a component of the deformation space is contractible if and only if $\ker I_* = \emptyset$. The whole deformation space is contractible if and only if $\ker I_* = \emptyset$, $G_0 = G_f$.

Lemma 4.1. *Let $\mathrm{Def}(f, A, B)$ be a 1-dimensional deformation space. Assume that γ_1 and γ_2 are disjoint arithmetically equalizing curves. Then $\mathrm{Def}(f, A, B)$ is not contractible.*

Proof. Assume the contrary. Each component M_0 of $M(f, A, B)$ is a Riemann surface with at least one puncture. Hence, $\pi_1(M_0)$ is a free group. If deformation space is contractible, then $\pi_1(M_0) = G_f$. There exists m_1, m_2 such that $T_{\gamma_1}^{m_1}, T_{\gamma_2}^{m_2} \in G_f$. This is a contradiction since $T_{\gamma_1}^{m_1}$ commutes with $T_{\gamma_2}^{m_2}$. □

Corollary 4.2. *Under the hypothesis of Lemma 4.1, if a component* Def_0 *of* $\mathrm{Def}(f, A, B)$ *is contractible, then* $\mathrm{Def}(f, A, B)$ *consists of countably many components.*

Proof. Let $M_0 = \pi(\mathrm{Def}_0)$. Since $\ker I_* = \emptyset$, $G_0 = \pi_1(M_0)$, G_0 is a free group subgroup of G_f. Since G_f contains two commuting elements of infinite order, G_0 has infinite index in G_f. Hence, $\mathrm{Def}(f, A, B)$ contains countably many components. □

5. Period 4 curve

Below we give a detailed description of $\mathrm{Per}_4 \subset \mathcal{M}_2^{cm}$ (compare [4]).

Using our standard parametrization, the every map in Per_4 takes the form

$$f_{b,c}(z) := z \mapsto 1 + \frac{b}{z} + \frac{c}{z^2}.$$

Let $\rho = f(1)$, then we have the following critical cycle: $0 \to \infty \to 1 \to \rho \to 0$.

$$1 + b + c = \rho$$
$$\rho^2 + b\rho + c = 0$$

$$b = \frac{-\rho^2 - \rho + 1}{\rho - 1}, \quad c = \frac{2\rho^2 - \rho}{\rho - 1}$$

Hence, the elements of Per_4 are parametrized by $\rho \in \hat{\mathbb{C}} \backslash \{0, 1, \infty, \frac{1}{2}\}$. We will denote by f_ρ the unique map in Per_4 with $f_\rho(1) = \rho$. The second critical point of f_ρ is

$$z_\rho = -\frac{2c}{b} = 2\frac{2\rho^2 - \rho}{-\rho^2 - \rho - 1}.$$

The corresponding critical value of f_ρ is

$$v_\rho = 1 - \frac{b^2}{4c}.$$

$$v_\rho = 1 - \frac{(\rho^2 + \rho - 1)^2}{4(\rho - 1)(2\rho^2 - \rho)}.$$

Fix a map $f_\rho \subset \mathrm{Per}_4$ such that $v_\rho \neq 0, 1, \rho, \infty$. Let $\mathrm{Per}_4' \subset \mathrm{Per}_4$,

$$\mathrm{Per}_4' := \{f_\rho \in \mathrm{Per}_4 : \ v_\rho \neq 0, 1, \rho\}.$$

$\mathrm{Per}_4 \setminus \mathrm{Per}_4'$ consists of 6 points. The map $I \colon \mathrm{Per}_4' \to \mathrm{Mod}(B)$

$$I(f_\rho) = (0, \infty, 1, \rho, v_\rho)$$

is an embedding, thus, Per_4' is a submanifold of $\mathrm{Mod}(B)$.

In this section we study the deformation space $\mathrm{Def}(f_\rho, A, B)$, $A = (0, 1, \infty, \rho)$, $B = (0, 1, \infty, \rho, v_\rho)$ for an arbitrary $f_\rho \in \mathrm{Per}_4'$.

Example of equalizing curves.

Consider a curve δ that separates 0 and v from ρ, 1 and ∞. All such curves are arithmetically equalizing, since $f^*(\delta) = i_*\delta = \emptyset$ (See figure 1)



FIGURE 1. Equalizing curve δ

We will show later that there are curves δ of this kind such that deformation space accumulates on $\mathcal{S}_\delta$, and there are curves δ such that deformation space does not accumulate on $\mathcal{S}_\delta$.

Lemma 5.1. *For $\rho \in \mathbb{R}_+$, let $\gamma_1 = \{\sqrt{\rho}e^{i\theta} : \theta \in [0, 2\pi]\}$. Then, for big enough ρ, $\gamma = f_\rho(\gamma_1)$ is an arithmetically equalizing curve and it separates $0, 1$ from v, ρ, ∞.*

Proof. Recall that

$$f_\rho(z) = 1 + \frac{b}{z} + \frac{c}{z^2},$$

$b = -\rho + O(1)$, $c = 2\rho + O(1)$. Therefore $f_\rho(\sqrt{\rho}e^{i\theta}) = -\sqrt{\rho}e^{-i\theta} + O(1)$. Since $v_\rho = -\frac{\rho}{8} + O(1)$ as $\rho \to \infty$, γ lies in a collar around β that does not contain $0, 1, \infty, \rho, v_\rho$. We see that γ_1 and γ are homotopic relative to B.

Let γ_2 be the second connected component of the preimage of γ; then γ_2 bounds unmarked preimages of 0 and 1, so it is homotopic to a point in $\hat{\mathbb{C}}\backslash B$. Hence γ is an arithmetically equalizing curve. □

Lemma 5.2. *The deformation space $\mathrm{Def}(f, A, B)$, where $f \in \mathrm{Per}_4'$ contains countably many components.*

Proof. Let δ be a curve that bounds ρ and v, and does not intersect γ.

We see that $[\delta]_A$ is peripheral as well as $[\delta_1]_A$ where $\delta_1 = f^{-1}(\delta)$ is the only connected component of the preimage of δ. We conclude that δ is arithmetically equalizing.

By Theorem 3.4 each component of Def is contractible. By corollary 4.2, the deformation space $\mathrm{Def}(f, A, B)$ contains countably many components. □

Even though δ is an equalizing curve, the deformation space $\mathrm{Def}(f, A, B)$ does not accumulate on $\mathcal{S}_\delta$.

Lemma 5.3. *The deformation space $\mathrm{Def}(f, A, B)$ does not accumulate on $\mathcal{S}_\delta$.*



FIGURE 2. A pair of equalizing curves γ and δ. Here and later: by x' we denote a point that has the same image as x.

Proof. Note that δ splits the sphere into two components, one with three marked points and a node and one with two marked points and a node. Since there is a unique conformal structure on the latter sphere $\mathcal{S}_\delta \cong \mathrm{Teich}(A)$. On the other hand, since δ_1 is peripheral $\mathcal{S}_{\delta_1} = \mathcal{S}_\emptyset = \mathrm{Teich}(A)$. We see that i_* restricts to $\mathrm{id}_{\mathrm{Teich}(A)}$ on $\mathcal{S}_\delta$.

Similarly, we see that the action of f^* on $\mathcal{S}_\delta$ restricts to the Thurston pull-back map $g^*\colon \mathrm{Teich}(A) \to \mathrm{Teich}(A)$ where g is a postcritically finite branched cover obtained from f by pinching δ_1 in the domain and δ in the range. The map g is obstructed: γ descends to a Levy cycle for g.

By continuity, any accumulation point τ of $\mathrm{Def}(f, A, B)$ satisfies $f^*(\tau) = i_*(\tau)$. Thus, any such accumulation on $\mathcal{S}_\delta$ would be a fixed point of g^*. Thurston's Theorem [1] guarantees that there are no fixed points for obstructed branched covers. □

6. Non-connectedness of the deformation spaces for maps in Per_n'

In this section we give an explicit construction of a pair of non-intersecting equalizing curves for maps in Per_n'. In view of Theorem 3.4 and Corollary 4.2, Lemma 6.2 yields the following result.

Theorem 6.1. *There exists $f \in \mathrm{Per}_n', n \geq 4$ such that $\mathrm{Def}(f, A, B)$ contains countably many components.*

Recall that with the parametrization where 0 is a critical point, $0 \to \infty \to 1$, a quadratic rational map takes the form

$$f_{b,c}(z) := 1 + \frac{b}{z} + \frac{c}{z^2},$$

We denote by $\rho_k(b, c) := f_{b,c}^{k-2}(1)$, $\rho_k = \frac{P_k(b,c)}{Q_k(b,c)}$, where $P_k(b, c)$ and $Q_k(b, c)$ are polynomials defined recursively by

$$P_2(b, c) = Q_2(b, c) = 1$$

$$P_{k+1}(b,c) = P_k^2(b,c) + bP_k(b,c)Q_k(b,c) + cQ_k^2(b,c),$$
$$Q_{k+1}(b,c) = P_k^2(b,c).$$

We see that for odd k, $\deg P_k = \deg Q_k + 1$ and for even k, $\deg P_k = \deg Q_k$.

The map $f_{b,c} \in \mathrm{Per}_n$ if and only if $\rho_n(b,c) = 0$.

Let p_k, q_k be the highest homogeneous part of P_k and Q_k correspondingly. Then for odd k,

$$p_{k+1} = p_k^2 + bp_kq_k, \quad q_{k+1} = p_k^2.$$

For even k,

$$p_{k+1} = q_k(bp_k + cq_k), \quad q_{k+1} = p_k^2$$

We will denote by $\tilde{p}_k$, $\tilde{q}_k$ reduced highest homogeneous parts of P_k and Q_k, obtained by canceling the common factors of p_k and q_k. Reduced homogenous parts satisfy the relation:

$$\tilde{p}_{k+1} = \tilde{p}_k + b\tilde{q}_k, \quad \tilde{q}_{k+1} = \tilde{p}_k \quad \text{for odd k;}$$
$$\tilde{p}_{k+1} = \tilde{q}_k(b\tilde{p}_k + c\tilde{q}_k), \quad \tilde{q}_{k+1} = \tilde{p}_k^2 \quad \text{for even k.}$$

Since $\tilde{p}_2 = 1$, $\tilde{q}_2 = 1$ do not have common factors, we can show inductively that $\tilde{p}_k$ and $\tilde{q}_k$, for $k \geq 2$ do not in fact have common factors.

The equation $\rho_n = 0$ defines an algebraic curve. The intersection of this curve with infinity in $\mathbb{CP}^2$ is given by solutions of the homogenous equation $\tilde{p}_n(b,c) = 0$.

Lemma 6.2. *Consider $f_{b,c} \in \mathrm{Per}_n'$ such that b, c are large, and $\frac{b}{c}$ is close to α. Let $\gamma_1 = \{\sqrt{|c|}e^{i\theta}, \theta \in [0, 2\pi]\}$. Then $\gamma = f_{b,c}(\gamma_1)$ separates $0, 1, \rho_4, \ldots, \rho_{n-2}$ from $v, \infty, \rho_3 \ldots, \rho_{n-1}$. Moreover, it is arithmetically equalizing.*

Proof. Note that if $\tilde{p}_k(b,c) = 0$ then $\tilde{p}_{km}(b,c) = 0$ for all m. Also, if $\tilde{p}_k(b,c) = 0$, then $\tilde{p}_{k+2m}(b,c) = 0$ and $\tilde{q}_{k+1+2m}(b,c) = 0$ for all m if k is odd.

We see that if n is odd, $\tilde{p}_n = \tilde{p}_{n-2}(b^2\tilde{q}_{n-2} + (b+c)\tilde{p}_{n-2})$ has a solution $(\alpha c, c)$ which is not a solution of $\tilde{p}_k$ for $k < n$. On the other hand if n is even, then any solution $(\alpha c, c)$ of $\tilde{p}_n$ is not a solution of $\tilde{p}_{2k+1}$ for $2k+1 < n$.

We observe that under the assumptions

$$f_{b,c}(\sqrt{|c|}e^{i\theta}) = \alpha\sqrt{|c|}e^{-i(\theta + \arg c/2)} + O(1).$$

Since α is not a solution of $\tilde{p}_k(\alpha c, c)$, $\deg p_k(\alpha c, c) > \deg q_k(\alpha c, c)$ for odd k, $\deg p_k(\alpha c, c) \leq \deg q_k(\alpha c, c)$ for even k. This yields $\rho_k(\alpha c, c) = \frac{p_k(\alpha c, c)}{q_k(\alpha c, c)} \geq a_k c + O(1)$ ($a_k \neq 0$) for odd k, $\rho_k(\alpha c, c) = \frac{p_k(\alpha c, c)}{q_k(\alpha c, c)} \leq b_k + O(1)$ ($b_k \neq 0$) for even k. Therefore, for big enough c, γ has the right separation property, and it lies in an annular neighborhood of γ_1 that does not contain any marked points.

Let γ_2 be the second preimage of the curve γ. For even n, γ_2 bounds unmarked preimages of $\infty, \rho_3, \ldots, \rho_{n-1}$. For odd n, γ_2 bounds ρ_{n-1} and unmarked preimages of $\infty, \rho_3, \ldots, \rho_{n-2}$. □

Lemma 6.3. *For any $n \geq 3$ there exists $f \in \mathrm{Per}_n'$, and a pair of non-intersecting equalizing curves γ and δ.*



FIGURE 3. A pair of equalizing curves for a map $f \in \mathrm{Per}_n$, n even



FIGURE 4. A pair of equalizing curves for a map $f \in \mathrm{Per}_n$, n odd

Proof. Take $f = f_{b,c}$ and γ from the previous lemma. Take a curve δ that encircles v and ρ_1 and does not intersect γ. □

7. Self-equivalences that preserve components of the deformation space

Let $\Gamma = \{\gamma_1, \dots, \gamma_n\}$ be a multicurve.

Lemma 7.1. *Consider a one-dimensional deformation space* $\mathrm{Def}(f, A, B)$. *Then Let* $\alpha : [0, 1) \to \mathrm{Def}(f, A, B)$ *be a path, such that* $\lim_{t\to 1} \gamma(t) \in \mathcal{S}_\Gamma$. *Then there exist* $m_1, \dots, m_n \in \mathbb{Z} \setminus \{0\}$ *such that* $T_{\gamma_1}^{m_1} \dots T_{\gamma_n}^{m_n} \in G_f$. *Moreover,* $T_{\gamma_1}^{m_1} \dots T_{\gamma_n}^{m_n}$ *preserves connected components of the deformation space.*

Proof. We first observe the following. Suppose that U is a holomorphic disk, $z \in U$, and $q\colon U \setminus \{z\} \to \mathrm{Mod}(B)$ is a holomorphic map that extends continuously (across z) to map U into the Deligne-Mumford compactification of $\mathrm{Mod}(B)$, and such that $q(z) \in \mathcal{S}_\gamma$ for some curve γ. Now let β be a curve in $U \setminus \{z\}$ that winds nontrivially around z; then $q(\beta)$ induces a nontrivial element of $\mathrm{Mod}(B)$, which in fact is T_γ^m for some $m \neq 0$. Similar arguments show that if $q(z) \in \mathcal{S}_\Gamma$ for some multicurve $\Gamma = \{\gamma_i\}$, then $q(\beta)$ induces a map of the form $\prod T_{\gamma_i}^{m_i}$ where all m_i are nonzero.

Now in the setting of the Lemma, $\mathcal{R}_{\mathrm{Rat}}(f, A, B)$ is a Riemann surface of finite type, and we must have $\pi_{\mathcal{TR}}(\alpha(t)) \to z$ as $t \to 1$, for some $z \in \overline{\mathcal{R}_{\mathrm{Rat}}(f, A, B)} \setminus \mathcal{R}_{\mathrm{Rat}}(f, A, B)$. Thus we can find a disk $U \subset \overline{\mathcal{R}_{\mathrm{Rat}}(f, A, B)}$ that maps into $\mathrm{Mod}(B)$ (by $\pi_{\mathcal{RM}}$) as described in the previous paragraph.

Then take any closed curve β in $\mathrm{Mod}(B)$ that is the image of a curve in $U \setminus \{z\}$ that winds once around z. We find that the image g_β of β in $\mathrm{MCG}(B)$ has the form $g_\beta = T_{\gamma_1}^{m_1} \dots T_{\gamma_n}^{m_n}$, where each $m_n \neq 0$. □

We give examples of paths on Per_4' that go around the punctures of Per_4' and corresponding equalizing curves. Recall that Per_4' is parametrized by ρ. The punctures are at points $\rho = 0, 1, \infty, 1/2$, as well as solutions of the equations $v_\rho = 0$, $v_\rho = 1$, $v_\rho = \rho$.

(1) Take a path that goes to a solution $v_\rho = \rho$. The path accumulates to $\mathcal{S}_\delta$, where δ is a curve that encircles v and ρ (see Figure 1).



FIGURE 5. Equalizing multicurve $\{\gamma_1, \gamma_2\}$

(2) Take a path that goes to a point $\rho = 0$, then $v \to \infty$. Hence, the path accumulates to $\mathcal{S}_\Gamma$, where $\Gamma = \{\gamma_1, \gamma_2\}$, γ_1 encircles ρ and 0, γ_2 encircles v and ∞ (see figure 5). $f^{-1}(\gamma_1) = \{\gamma_1^1, \gamma_1^2\}$, $f^{-1}(\gamma_2) = \{\gamma_2^1, \gamma_2^2\}$, γ_1^1, γ_1^2 and γ_2^1 are peripheral. $f^*(\gamma_1) = \emptyset$, $f^*(\gamma_2) = \gamma_2^2$, $i_*(\gamma_1) = \gamma_2^2$, $i_*(\gamma_2) = \emptyset$ (see figure 5)

(3) Let α be a path from a given point to $\rho = \infty$. We calculate $v_\rho \sim -\frac{\rho}{8}$. Hence, the path accumulates to $\mathcal{S}_\gamma$, γ is a curve that encircles v, ρ, ∞ (see figure 1). Thus, T_γ preserves components of the deformation space. Let δ

be a curve that encircles v and ρ and does not intersect γ (see figure 1). Since, T_δ^2 commutes with T_γ, the map T_δ^2 permutes the components.

(4) Similarly, if we take f and γ as in Lemma 6.2, then for odd n, T_γ preserves components of the deformation space. Let δ be a curve that encircles v and ρ, and does not intersect γ (see figures 4, 3). Since, T_δ^2 commutes with T_γ, the map T_δ^2 permutes the components.

References

[1] A Douady and J.H. Hubbard, *A proof of Thurston's characterization of rational functions.*, Acta Math. **171** (1993), no. 2, 263-297.

[2] A.L. Epstein, *Transversality in Holomorphic Dynamics.*, preprint, http://homepages.warwick.ac.uk/∼mases/Transversality.pdf.

[3] E. Hironaka and S. Koch, *A disconnected deformation space of rational maps*, preprint.

[4] J. Milnor, *Geometry and Dynamics of Quadratic Rational Maps*, Experimental Mathematics **2**, 37-83.

[5] M. Rees, *View of Parameter Space: Topographer and Resident.*, Astrisque **288** (2003).

[6] S. Koch, K. Pilgrim, and N. Selinger, *Pullback invariants of Thurston maps*, Transactions of the American Mathematical Society (to appear).

[7] N. Selinger, *Thurstons pullback map on the augmented Teichmuller space and applications*, Invent. Math. **189** (2012).